

HYPOGLYCAEMIA

LOW if Blood Glucose Level is below:
TREAT IMMEDIATELY

Signs and symptoms

Note: Symptoms may not always be obvious

**DO NOT LEAVE STUDENT UNATTENDED
DO NOT DELAY TREATMENT**

Student conscious & cooperative

Able to eat hypo food

Hypo treatment or fast acting carb:

As supplied or listed on management plan

Recheck BGL after 15 mins

If BGL repeat fast acting carb

Student unconscious or drowsy

Risk of choking or unable to swallow

First aid

Place student on their side and stay with the student

**CALL AN AMBULANCE
DIAL 000**

Contact parent or guardian

when safe to do so

To be used in conjunction with management plan

Student's name

DOB

Grade/Year

Key contacts

Phone number

Phone number

Injection

Will an injection will be required at school:

no	yes
	with supervision
	by a staff member

In room/location

Continuous Glucose Monitoring (CGM)

Routine BGL checking times

- Anytime, anywhere in the school
- Prior to lunch and other times as per management plan
- Any time hypo is suspected or student feels unwell
- Prior to activity
- Prior to exams or tests

HYPERGLYCAEMIA

HIGH if Blood Glucose Level is above
(High BGLs are not uncommon)

Signs and symptoms

Note: Symptoms may not always be obvious

Student well

Re-check BGL in 2 hours

Encourage student to drink water and return to class

In 2 hours, if BGL still above contacts for advice

Student unwell

e.g. vomiting

Check ketones

(Refer to plan)

Call key contact(s) to collect student ASAP

call key

Clinic:

Clinic contact:

Contact No: